

A Non-singular Theory of Gravity

Neil J. Cornish and John W. Moffat
Department of Physics, University of Toronto
Toronto, Ontario M5S 1A7, Canada

Abstract

We present a geometrical gravitational theory which reduces to Einstein's theory for weak gravitational potentials and which has a singularity-free analog of the Schwarzschild metric.

Despite its geometrical elegance, Einstein's General Theory [1] is plagued by singular solutions which destroy global predictability. It has long been hoped that a new, non-singular theory of gravity might be found, from which Einstein's theory emerges as a low-energy limit. We shall present the static spherically symmetric solution to a close relative of Einstein gravity, which suggests that we may have found such a theory.

The essential new physics comes from endowing the gravitational field with additional degrees of freedom. This additional freedom gives the field the ability to smooth out the energy density of a point source over a small but finite region, thus removing any singular behaviour.

The theory we shall study was first put forward as a theory of gravity in 1979 [2], after it was realized that the field equations of Einstein-Straus Unified Field Theory [3] did not describe a unification of electromagnetism and gravity, but rather a generalized theory of gravity. Unlike General Relativity, the metric tensor in this theory is not assumed to be symmetric. This means that both the manifold and the tangent space transform under $GL(4, R)$, while in General Relativity the tangent space is restricted to the Lorentz subgroup, $SO(3, 1)$, of $GL(4, R)$. The resulting theory was called Nonsymmetric Gravitational Theory (NGT).

The phenomenology of NGT which has been described in the literature to-date [4] turns out to be a singular subset of the full theory. Both General Relativity and the previously studied sector of NGT emerge as singular limits of the complete theory.

We shall present an exact, non-singular analog of the Schwarzschild solution for NGT. The spacetime this solution describes has no event horizons and no curvature singularities. The solution set is parameterized by two constants of integration, m and s . We can identify m as the gravitational mass by taking the Newtonian limit of the solution. The second constant, s , is a dimensionless coupling constant which we shall call the skewness parameter, since it controls the size of the skew-metric contributions. We shall see that in the weak field limit, the skew sector of the theory behaves as sm^2/r^2 for large r . The non-linear coupling to mass emphasizes the purely geometrical nature of this new sector.

For strong fields we shall see that the limit $s \rightarrow 0$ is non-analytic. This means that there are no event horizons or singularities for any $s > 0$, while there are both horizons and curvature singularities when $s = 0$. That is what we mean when we say that General Relativity (and the old solutions to NGT) are singular limits of the full theory.

The most general solution to the field equations for NGT introduces a third constant of integration, l^2 . This constant is known as the NGT charge, and has the dimensions of a length squared. Since this charge arises from a conserved current, it has been postulated that it is proportional to conserved particle number. All previous work on NGT has been based on this charge being non-zero. To simplify our current presentation, we shall take the NGT charge to be zero. This results in no essential loss of generality since the full theory is non-singular for arbitrary l^2 , so long as $s > 0$. Moreover, this choice emphasizes the fact that no new matter couplings are needed to obtain non-singular solutions.

The NGT Lagrangian without sources is given by [4]

$$\mathcal{L} = \sqrt{-g}g^{\mu\nu} \left(R_{\mu\nu}(\Gamma) + \frac{2}{3}W_{[\mu,\nu]} \right) , \quad (1)$$

where $R_{\mu\nu}(\Gamma)$ is the generalized Ricci tensor, W_μ is a Lagrange multiplier and Γ refers to the

torsion-free connection $\Gamma_{\mu\nu}^\lambda$. Square brackets denote anti-symmetrization and we use units where $G = c = 1$ throughout. The vacuum field equations that follow from (1) are

$$g_{\mu\nu,\sigma} - g_{\rho\nu}\Gamma_{\mu\sigma}^\rho - g_{\mu\rho}\Gamma_{\sigma\nu}^\rho = 0, \quad (2)$$

$$(\sqrt{-g}g^{[\mu\nu]})_{,\nu} = 0, \quad (3)$$

$$R_{\mu\nu}(\Gamma) = \frac{2}{3}W_{[\nu,\mu]}. \quad (4)$$

The most general, spherically symmetric metric for NGT was found by Papapetrou [5] to be

$$g_{\mu\nu} = \begin{pmatrix} \gamma(r) & w(r) & 0 & 0 \\ -w(r) & -\alpha(r) & 0 & 0 \\ 0 & 0 & -r^2 & f(r)\sin\theta \\ 0 & 0 & -f(r)\sin\theta & -r^2\sin^2\theta \end{pmatrix}. \quad (5)$$

In this paper we shall take the NGT charge to be zero which demands $w(r) = 0$. Papapetrou gave a special, unphysical solution to field equations for this metric which had $f(r) = Ar^2$. This solution is discarded since the resulting metric is not of Minkowskian form for large r . While studying the general wave behaviour of NGT [6], we found that asymptotically Minkowskian solutions could be found with $f(r) = \text{const.}$, since this corresponds to $1/r^2$ fall-off in orthonormal coordinates. The static spherically symmetric solutions found by Wyman [7], Bonnor [8] and Vanstone [9] for the field equations of Unified Field Theory share this asymptotic behaviour [10]. This lead us to adapt their solutions to NGT, and we found that the resulting solution proved to be free of singularities.

The metric functions are given by

$$\gamma = e^\nu, \quad (6)$$

$$\alpha = \frac{m^2(\nu')^2 e^{-\nu}(1+s^2)}{(\cosh(a\nu) - \cos(b\nu))^2}, \quad (7)$$

$$f = \frac{2m^2 e^{-\nu}(\sinh(a\nu)\sin(b\nu) + s(1 - \cosh(a\nu)\cos(b\nu)))}{(\cosh(a\nu) - \cos(b\nu))^2}, \quad (8)$$

where

$$a = \sqrt{\frac{\sqrt{1+s^2}+1}{2}}, \quad b = \sqrt{\frac{\sqrt{1+s^2}-1}{2}}, \quad (9)$$

prime denotes differentiation with respect to r , and ν is given implicitly by the relation:

$$e^\nu(\cosh(a\nu) - \cos(b\nu))^2 \frac{r^2}{2m^2} = \cosh(a\nu)\cos(b\nu) - 1 + s\sinh(a\nu)\sin(b\nu). \quad (10)$$

For small r this relation has multiple branches of solutions for ν , but only one of these branches matches onto the unique solution of ν for large r . We shall only be interested in the unique inversion of (10) which yields an asymptotically flat spacetime.

Unfortunately, we can only invert (10) analytically for $r/m \ll 1$ and $m/r \ll 1$ and we must resort to numerical methods to establish the intermediate behavior.

Starting in familiar territory, we find for $m/r \ll 1$ and $s \ll 1$ that the metric takes the near-Schwarzschild form:

$$\gamma = 1 - \frac{2m}{r} + \frac{s^2 m^5}{15r^5} + \frac{4s^2 m^6}{15r^6} + \dots, \quad (11)$$

$$\alpha = 1 + \frac{2m}{r} + \frac{4m^2}{r^2} + \frac{8m^3}{r^3} + \frac{(16 - 2s^2/9)m^4}{r^4} + \dots, \quad (12)$$

$$f = \frac{sm^2}{3} + \frac{2sm^3}{3r} + \frac{6sm^4}{5r^2} + \dots, \quad (13)$$

where the higher order terms in m/r include higher powers of s also. We see that for large r the NGT corrections to the Schwarzschild solution are small for any $s < 1$, and can be arbitrarily small if s is close to zero. Clearly, experimental predictions such as the bending of light by the sun and the perihelion advance of Mercury will only be minutely affected by having s non-zero.

Near the origin we can develop expansions where $r/m \ll 1$ and $0 < s \ll 1$. The leading terms are

$$\gamma = \gamma_0 + \frac{\gamma_0(1 + \mathcal{O}(s^2))}{2s} \left(\frac{r}{m}\right)^2 + \mathcal{O}((r/m)^4), \quad (14)$$

$$\alpha = \frac{4\gamma_0 + \mathcal{O}(s^2)}{s^2} \left(\frac{r}{m}\right)^2 + \mathcal{O}((r/m)^4), \quad (15)$$

$$f = m^2 \left(4 - \frac{s\pi}{2} + s^2 + \mathcal{O}(s^3)\right) + \frac{s + s^2\pi/8 + \mathcal{O}(s^3)}{4} r^2 + \mathcal{O}(r^4), \quad (16)$$

where γ_0 is given by

$$\gamma_0 = \exp\left(-\frac{\pi}{s} - 2 + \mathcal{O}(s)\right). \quad (17)$$

The above equation clearly illustrates the non-analytic nature of the limit $s \rightarrow 0$ in the strong field region. We shall postulate that s is always strictly greater than zero, and refer to this new phenomenology for NGT as Non-Singular Gravity or NSG.

Our numerical results confirm that all the expansions we have exhibited produce excellent approximations to the exact solution in their respective regions of validity.




FIG. 1. $\gamma(r)$ in the region around $r = 2m$ for both GR ($s = 0$) and NSG with the choice $s = 0.1$.

Figure 1. shows how γ behaves in the region where neither of our expansions are valid, for the choices $s = 0$ (GR) and $s = 0.1$. We see that for NSG the redshift is finite between any two points in the spacetime. The maximum redshift is between $r = 0$ and $r = \infty$, and is given by

$$z = \exp\left(\frac{\pi}{2s} + 1 + \mathcal{O}(s)\right) - 1. \quad (18)$$

Put another way, we find that a timelike Killing vector at spatial infinity remains timelike throughout the spacetime - our solution is free of event horizons.




FIG. 2. The acceleration of an initially static observer in the region near $r = 2m$ for NSG with $s = 0.1$ and $m = 1$.

Figure 2. shows the inward radial acceleration of a static observer near $r = 2m$ for the choice $s = 0.1$. Unlike GR, we see that the acceleration is finite at $r = 2m$. The radial

acceleration in NGT is given by the familiar expression $a^r = 1/2(\ln \gamma)' = \nu'/2$. Near $r = 0$ we find

$$a^r = \frac{1 + \mathcal{O}(s^2)}{2sm^2}r + \mathcal{O}(r^3/m^4) , \quad (19)$$

while for large r we have

$$a^r = \frac{m}{r^2} + \frac{2m^2}{r^3} + \frac{4m^3}{r^4} + \frac{8m^4}{r^5} + \frac{(16 - s^2/6)m^5}{r^6} + \dots . \quad (20)$$

At large distances we have a new, repulsive, $1/r^6$ contribution to the gravitational “force” which is of little interest. In contrast, the situation near $r = 0$ is dramatically different from GR, since the “force” vanishes in NSG whereas it is infinite in GR. The shape of the acceleration curve for initially static observers in NSG is reminiscent of the acceleration curves one finds for extended matter distributions such as planets or stars. The acceleration goes roughly as $1/r^2$ outside the body and tails off smoothly to zero inside the body. We attribute this to the fact that in NSG the degrees of freedom associated with the skew field f are able to distribute the energy of the point source over an extended region. Figure 3. shows a plot of the invariant $F = (g_{[\theta\phi]}g^{[\theta\phi]})^{1/2} = f/\sqrt{f^2 + r^4}$. For large r we find $F \sim sm^2/3r^2$, while at the origin $F = 1$.




FIG. 3. The invariant skew potential near $r=2m$ for NSG with the choice $s = 0.1$

We see that F is negligible outside of $r = 2m$, while below $r = 2m$ it quickly rises to 1. The radial dependence exhibited by F is similar to the energy density profile of a compact object with radius $r = 2m$, and suggests the mechanism by which the singularities can be avoided.

The vanishing of the radial acceleration at $r = 0$ leads us to expect that the curvature is non-singular at the origin. This expectation is confirmed by an explicit calculation of the generalized curvature invariants. NGT generally has more curvature invariants than GR. However, the fact that $R_{[\mu\nu]} = 0$ for the solution we are studying reduces the number that

needs to be calculated to four. Direct numerical evaluation shows that the curvature is non-singular for all r . Since there is a coordinate singularity in α at $r = 0$, it is also important to establish analytically that there is no curvature singularity at $r = 0$, rather than relying on our numerical results. Using the expansions (14), (15) and (16), we found all the curvature invariants to be finite. A representative invariant is the generalized Kretschmann scalar, which is given to leading order in s by

$$K = R^{\mu\nu\kappa\lambda} R_{\mu\nu\kappa\lambda} \quad (21)$$

$$= \frac{\exp(\pi/s + 2)}{2sm^4} , \quad \text{at } r = 0 , \quad (22)$$

$$= \frac{48m^2}{r^6} , \quad r \rightarrow \infty . \quad (23)$$

All the non-vanishing curvatures are of the form $1/m^2$ at the origin and m/r^3 for large r , which is to be expected on dimensional grounds. These results are again reminiscent of some extended energy distribution. For example, the non-vanishing curvature invariants at the center of a star are proportional to the local energy density ρ . Consistent with this picture we see that the curvature is smaller for larger values of s since for large s the skew field is better able to smooth out the would-be singularity at $r = 0$.

While all the results we have presented suggest that NSG sector of NGT is indeed a non-singular theory of gravity, there is still much to be understood. An obvious question is whether the spacetime we have described would ever be reached as the endpoint of gravitational collapse. Since gravity tends to turn off at very short distances, it is perhaps more likely that some kind of super-dense object might form with the metric we have described as its exterior spacetime. Observationally it would be difficult to distinguish such objects from black-holes. Looking beyond the limitations of a static spherically symmetric solution, we need to understand the role played by the s parameter in more general settings. Is s related to a new fundamental constant which controls the cross-coupling of the skew and symmetric parts of the geometry, or is it just a parameter which varies from one solution to the next? At the linearized level, we know [6] that s can be considered as a coupling constant between the three divergence-free parts of $g_{[\mu\nu]}$ and the rest of the $g_{\mu\nu}$'s, but this understanding is of little use in the strong field regime where these degrees of freedom are most important. Another interesting question is whether a similar smoothing effect by the skew metric can eliminate cosmological singularities such as the Big Bang. For now we will have to be content with a non-singular analog of the Schwarzschild geometry.

ACKNOWLEDGEMENTS

This work was supported by the Natural Sciences and Engineering Research Council of Canada. One of the authors (NJC) is grateful for the support provided by a Canadian Commonwealth Scholarship. We thank P. Savaria, M. Clayton, B. Holdom and J. Levin for helpful discussions.

REFERENCES

- [1] Einstein A., *Ann. der Physik*, **49**, 1916.
- [2] Moffat J. W., *Phys. Rev. D* **19**, 3554, 1979; *Phys. Rev. D* **35**, 3733, 1987.
- [3] Einstein A. and Straus E. G., *Ann. Math.* **47**, 731, 1946.
- [4] For a review of early work on NGT see Moffat J. W. in *Gravitation - A Banff Summer Institute*, eds. Mann R. B. & Wesson P., World Scientific, Singapore, p. 523, 1991.
- [5] Papapetrou A., *Proc. Roy. Irish Acad. Sci.* **A52**, 69, 1948.
- [6] Cornish N. J., Moffat J. W., *University of Toronto preprint* UTPT-94-05, 1994.
- [7] Wyman M., *Can. J. Math.* **2**, 427, 1950.
- [8] Bonnor W. B., *Proc. Roy. Soc.* **209**, 353, 1951; **210**, 427, 1952.
- [9] Vanstone J. R., *Can. J. Math.* **14**, 568, 1962.
- [10] Savaria P., *private communication*.